

Calf immunity – feeding during the critical first two weeks

by Julia Hamann, Ruminant Marketing Manager, Diamond V, Cedar Rapids, Iowa, USA.

During the first two weeks after birth, a calf's health and future productivity is at high risk. Investing time and money to reduce the incidence of disease and improve survival at this time generates good payback. Feeding to support the calf's immune system, combined with good husbandry and environment, helps to optimise overall health, growth, and productivity.

The newborn calf possesses a functional immune system, but it is poorly developed. Antibodies such as immunoglobulins (IgG) do not cross the placenta in pregnancy. However, they are available to the calf through colostrum. In order to help protect from diseases early in life, the calf depends on receiving adequate amounts of high quality colostrum soon after birth.

Feeding good colostrum offers the first opportunity to support the calf's immune system. Practical advice includes the 'Five Qs' – a memory aid for colostrum management (see Fig. 1).

As the calf's immune system develops, it is important to keep the gut healthy. Roughly 70% of all immune cells are part of the gastro-intestinal tract in the form of 'gut associated lymphoid tissue' (GALT).

GALT is the calf's first line of defence and a potential entry site for disease-causing pathogens and infectious agents. Any disruption of GALT impacts the calf's immune system.

Disease susceptibility

Although the calf's gastro-intestinal tract is developing rapidly, the acquisition of immunity requires time. During the first few weeks after birth, the calf remains susceptible to diseases.

The most common cause of death in pre-weaned calves results from gastro-intestinal pathogens, such as *Escherichia coli*, salmonella, and *Clostridium* bacteria, *Cryptosporidia* protozoa, and Rotavirus-Coronavirus virus.

In 2007, a study of the US Department of



Agriculture's National Animal Health Monitoring System (USDA NAHMS) reported an average 7.8% mortality rate in pre-weaned dairy heifers. Digestive problems accounted for 56.5% of the deaths.

In a more recent US survey of 105 calf ranches, mortality averaged 3.6% at operation level (average of total number of farms surveyed) and 5.4% at calf level (average of total number of calves in the survey). In this survey, more than half of the deaths were attributed to diarrhoea.

In other parts of the world, similar mortality rates occur. In Quebec, Canada, a study involving 115 farms reported a pre-weaned calf mortality of about 8%.

In Europe, 60 farms in Austria and

Germany showed a mortality rate of about 5%. Therefore, improvements in gut health to minimise susceptibility to disease are likely to help to minimise mortality in pre-weaned calves.

Feeding strategies

Feeding strategies for the very young calf have been changing. In 2010, 7-10% of pre-weaned heifers affected by digestive problems were treated with antibiotics.

Today, 'all-natural' feeding strategies have become important in helping to maintain calf health and productivity, including during the critical first two weeks after birth.

These feeding strategies have focused on decreasing the interaction of potential pathogens with cells in the gastro-intestinal tract.

Feed additive products used in these strategies have included hyper-immunised egg protein, probiotics, prebiotics, and microbial fermentation metabolites.

Various studies have shown beneficial effects from supporting gastro-intestinal health and growth in pre-weaned calves.

Hyper-immunised egg proteins are the product of eggs inoculated with specific pathogenic organisms in order to produce specific antibodies to those organisms. These proteins are then fed to the calf to help minimise gastro-intestinal diseases.

Probiotics or 'direct-fed microbials' are live bacteria that are typically combinations of various species. They may include combinations of benign lactic acid producing

Continued on page 27

Fig. 1. The five 'Qs' of colostrum management (S. Godden, 2013).

- Quality: 50g/L IgG.
- Quantity: 10% body weight (roughly four litres for a normal dairy calf).
- Quickness: 1-2 hours after birth.
- SQueaky clean: Low bacterial contamination (TCC [total coliform count] <10,000 CFU/ml; TPC [total plate count] <100,000 CFU/ml).
- Quality of passive transfer: More than 80% of calves should have TP (total protein) \geq 5.5g/dl.

Continued from page 25

species that can propagate in the intestines to help prevent attachment and colonisation by pathogenic bacteria.

However, because these are live bacteria, some probiotic products require coating or other treatments in order to survive and remain effective following storage, feed processing heat and pressure, and the pH of the gastro-intestinal tract.

Prebiotics are dietary components that are not digested by the calf, which promote growth of beneficial bacteria in the calf's gastro-intestinal tract.

The growth of beneficial bacteria help maintain and possibly improve overall gut health and digestibility.

In 2011, Ballou conducted a study to determine the effects of supplementation of a blend of probiotics, prebiotics, and hyper-immunised egg proteins in Holstein calves during the three first weeks after birth.

Results from this study showed an improvement in gastro-intestinal health. Various other studies have also shown beneficial effects of these products in supporting gastro-intestinal health, as well as growth in pre-weaned calves. However, results from studies on the benefits of probiotics and prebiotics on health of dairy calves remain ambiguous.

Some studies have shown a reduction in scouring and improvement in growth, while other studies have shown no benefits of

including either prebiotics or probiotics in milk for calves. The lack of a clear effect in calves is likely due to many factors – such as environment, age of the calf, rearing techniques, etc – which require additional investigation.

Fermentation products

Liquid feeding strategies have become popular to deliver some of these gut health products to the calf during the critical first two weeks of life when the calf does not eat much dry feed.

Recent research sponsored by Diamond V used this approach to deliver a new nutritional health product called SmartCare to newborn calves. This product is composed of unique bioactive metabolite compounds produced during precision fermentation of *Saccharomyces cerevisiae*.

A study with this product in the milk replacer and Original XPC in the starter ration was conducted by Iowa State University with pre-weaned dairy calves inoculated with live salmonella bacteria.

A second study conducted by researchers at Texas Tech University and the USDA ARS validated the Iowa State results and provided additional data on calf starter intake, average daily gain, and feed efficiency when salmonella challenged calves were fed these products compared to a placebo control group.

Higher plane of nutrition

When following a liquid feeding strategy to support calf health and productivity in the first two weeks after birth, there are many options. Not all probiotics, prebiotics, hyper-immunised egg proteins, or fermentation products perform alike.

As a rule, those based on peer-reviewed controlled research are more likely to achieve predicted outcomes.

After the first two weeks, the calf's risk of gastro-intestinal diseases decreases dramatically. Also, the calf's plane of nutrition, achieved during the first two weeks and for the remainder of the pre-weaning period, has effects on the calf's immune responses. These effects extend into the immediate post-weaning period.

A recent meta-analysis suggests that higher planes of calf nutrition positively impact the long-term productivity of the heifer and cow.

Developing new feeding strategies to support immune function in the newborn calf is an exciting area for research requiring more studies. These strategies can help optimise health and growth during the critical first two weeks and support long term productivity in the heifer and cow.

*References are available on request
jhamann@diamondv.com*