

Impact of milk fever on the lifetime performance of your dairy cows

Milk fever is a metabolic disturbance or production disease of dairy cows that generally occurs just before, or soon after, calving due to low calcium levels in the blood. It is associated with the drain of calcium within the foetus and milk during pregnancy and calving, respectively. Milk fever has both direct and indirect economic impacts on the dairy industry.

by **Dr Maurice Verschaeve, Veterinary Surgeon,**
Kathleen Verschaeve, Pharmacist,
Jonas De Roo, Biologist &
Dr Glenn Verstraete, Pharmacist,
Emma Nutrition.
www.emma.be

The most important direct economic losses due to milk fever are losses due to reduction in milk production of affected cows, loss of animals through death and culling, and the cost of treatment.

In addition, the excessive drop in calcium levels around calving can



Single dose Green Bolus that can slowly release calcitriol-glycoside.

lead to a long-term calcium deficiency, further threatening the economic profitability. Hence, the prevention of milk fever and long-term calcium deficiency is key in the dairy industry.

Milk fever

Milk fever (also called parturient paresis or parturient hypocalcaemia) is one of the production diseases which primarily occur among older high-yielding dairy cows during the period around calving. The increased demand for calcium at parturition due to the onset of lactation may result in a calcium deficiency situation potentially provoking the outbreak of milk fever in dairy cows.

Milk fever generally occurs when calcium blood plasma concentration is lower than 8mg/dL (i.e., 2.2 mmol/L). Such a low calcium blood plasma concentration has consequences for many nerve and muscle functions.

Milk fever is an economically important disease due to milk secretion that can reduce the productive life of a dairy cow. Milk fever incidence has remained steady in some countries at about 10%.

Milk fever affects about 6% of dairy cows in the United States each year, according to the 1996 National Animal Health Monitoring Survey. If left untreated, about 60-70% of the affected cows may die.

There are some factors which influence incidence and severity of milk fever. These are: age (older cows are more sensitive than younger),



milk yield (cows with higher yield are more predisposed than cows with lower yield), breed, body condition, length of dry period and diet composition.

Long-term hypocalcaemia

As mentioned earlier, high yielding dairy cows suddenly experience a high demand of calcium during parturition.

Calcium is necessary for contraction of the uterine wall muscle and milk production. The natural regulation systems in the dairy cow can not always cope with this sudden increase in calcium demand. As a result, calcium blood plasma levels can drop significantly.

Depending on the severity of this drop in calcium levels this can lead to a long-term calcium deficiency and thus a persistent lack of appetite, hypersensitivity, weakness and, if left untreated, even paralysis and mortality.

Long-term hypocalcaemia also has some widespread effects that predispose to other periparturient diseases such as mastitis, ketosis, displaced abomasum and retained placenta. Even if no visible symptoms are observed, long-term hypocalcaemia can lead to a decrease in production and fertility.

Even successful treatment of hypocalcaemia does not eliminate further complications associated with milk fever, which results in further economic losses. This stresses the demand for an effective prevention: administration of calcitriol.

The role of calcitriol

Calcitriol or 1,25-dihydroxyvitamin D3 is a metabolite of vitamin D3. Unlike vitamin D3 that needs to be transformed in the liver and kidney, calcitriol works directly on the vitamin D3 receptors in the intestinal wall, bones and kidney having the following effects:

- **Intestinal wall:** Calcitriol is able to activate the calcium-binding proteins increasing the calcium absorption rate from the gut to the blood vessels.

- **Bones:** Calcitriol activates the vitamin D receptors in the bone cells increasing calcium deposition or release from the bones, depending on the concentration of calcium in the blood plasma.

- **Kidney:** Calcitriol stimulates the resorption of calcium from the urinary tracts to the blood vessels


Taking all the above-mentioned into account, it can be stated that calcitriol can increase the absorption and regulation rate of calcium thus maintaining an optimal calcium level in the blood plasma. In addition, calcitriol is known to have a similar effect on phosphorus absorption and regulation, positively affecting the immune system and improving the suppression of pathogens.

Advantages of calcitriol-glycoside

The binding of calcitriol to a glycoside is known to improve

Continued on page 14

Fig. 1. Impact of gastrointestinal residence time (days) on bolus weight (g).



Continued from page 13 stability and increase bioavailability (due to better absorption through the intestinal wall). Calcitriol-glycoside can be found in dried leaves of the plant *Solanum glaucophyllum* (containing a standardised concentration).

Studies with herbal calcitriol-glycoside added to the feed of ruminants show a significant increase of calcitriol in the blood levels. Furthermore, it was observed that a single high dose of herbal calcitriol-glycoside around calving could result in a major increase in both calcium and phosphorus blood levels.

Slow-release calcitriol-glycoside

Considering calcitriol-glycoside as an effective substance to prevent hypocalcaemia, it was of the highest importance for Emma Nutrition to search for the most convenient and easiest way of administration.

As displayed in Fig. 1, Emma Nutrition developed a slowly

disintegrating bolus that should only be administered once (approximately 1-2 days before calving).

As listed in Table 1 and displayed in Fig. 2, this unique patented formula allows farmers to easily increase calcitriol blood concentration 8-10 hours after oral administration.

Furthermore, it was found that calcium concentration in blood could be kept above 2.2 mmol/L (i.e., 8.8mg/dL which is the recommended minimum calcium blood plasma concentration to prevent milk fever and long-term hypocalcaemia).

Therefore, one might assume that hypocalcaemia could still be countered even in case the Green Bolus is administered just before or after calving. Finally, it was observed that the administration of Green Bolus could positively affect phosphorus levels. As a result of the excellent in vivo performance, Emma Nutrition's Green Bolus was officially recognised in 2020 as a highly effective diet food for the prevention of long-term hypocalcaemia and milk fever.

Calcium concentration in blood (mmol/L)			
	Cow	T0 (when applying bolus)	T1 (24 hours after calving)
Control	1	2.33	2.13
	2	1.93	1.83
			(2.93 - 24 hours after Green Bolus)
Green Bolus	3	1.87	2.47
	4	1.98	2.16
	5	1.96	2.16
	6	2.51	2.16
	7	2.10	2.21
	8	2.20	2.25
	9	2.24	2.33

Table 1. Impact of Green Bolus administration on calcium concentration in the blood of Holstein-Friesian cattle.

Conclusion

Emma Nutrition successfully developed a single dose calcitriol glycoside bolus (Green Bolus) that is officially recognised as a diet food to prevent dairy cows from

hypocalcaemia and milk fever, resulting in a higher economic yield.

References are available from the authors on request

Fig. 2. The impact of Green Bolus administration on cholecalciferol (left) and calcium, phosphorus and magnesium levels (right).

