

# Isoleucine and its role in crude protein reduction in broilers

After the incorporation of the major five amino acids namely lysine, methionine, threonine, valine and arginine, the commercial nutritionists are experiencing the pressure point on L-isoleucine to meet its requirements. Generally, the L-isoleucine requirement is either being ignored or compromised due to its non-availability commercially.

by Dr Daulat Khan,  
CJ Europe GmbH, Germany.  
www.cjbio.net

Now, it has recently become commercially available, thus it is important to define its current status in the commercial feeds as well as identify the compromises on its requirement in order to achieve the best performance with controlled economics.

## Requirements of L-isoleucine

Recommendations from breeding companies for the L-Isoleucine requirements and the literature studies are summarised in Tables 1 and 2.

The average trend is showing an increase in dlle:dLys with the age period.

According to recommended standards the average dlle:dLys for the starter, grower and finisher periods is 0.67, 0.68 and 0.69, respectively, whereas, an individual data available based on different age periods and genetics also shows variable results (Table 2).

Contrary to this, commercial poultry diets are not meeting the

**Table 1. Ideal digestible isoleucine to digestible lysine (dlle:dLys) ratio recommendations.**

Reference	dlle:dLys		
	Starter	Grower	Finisher
Ross 308, 2019 (Also all veg diet)	0.67	0.68	0.69
Cobb 500, 2018	0.63	0.64	0.66
Brazilian, 2017	0.67	0.68	0.68

Experimental period (days)	Estimated dlle requirement (%)	Strain	Reference
7-21	0.68	Cobb 500 x Cobb 500	Helmbrecht et al., 2010
7-21	0.72	Cobb 500 x Cobb 500	Campos et al., 2009
18-30	0.62	Ross 308 x Ross 308	Kidd et al., 2004
22-42	0.65	Ross x Hubbard	Kidd et al., 2000
28-40	0.69	Cobb 500 x Cobb 500	Campos et al., 2009
28-42	0.65	Ross 708 x Ross 708	Mejia et al., 2011
30-42	0.59	Ross 308 x Ross 308	Kidd et al., 2004
30-43	0.66	Cobb 500 x Cobb 500	Helmbrecht et al., 2010
42-56	0.55	Ross 308 x Ross 308	Kidd et al., 2004
22-42	0.72	Cobb 500	Duarte et al., 2015

**Table 2. Individual studies on the recommendations of dlle.**

recommended level of L-isoleucine, which is even challenging in low crude protein (CP) diets. The estimated Ile to Lys ratio in commercial diets can vary from 50-66:100. These variations are related to age period, raw material, and the formulation limits or even no limit check on Ile during the formulation.

## Balance between branched chain amino acids (BCAAs) in raw materials

Raw materials contain enough of leucine (Leu) as compared to Ile and Valine (Val). Table 3 summarises the ratios between Ile:Lys, Leu:Lys and Val:Lys in commonly used raw materials.

The antagonism of the BCAA is described as early as 1955, which can decrease feed intake or cause leg weakness and torn feathers in growing chickens.

Therefore, an imbalance between BCAAs should be avoided in order to achieve the most optimum performance.

Studies are also available which do not show any performance difference due to higher Leu.

Waldroup et al., (2002) concluded that the antagonism among or between BCAAs is not likely to reduce the performance of broilers under practical conditions.

Erwan et al., (2009) suggested that increasing dietary leucine had no significant effect on feed consumption, weight gain and feed to gain ratio and carcass characteristics, but significantly increased carcass weight up to 9% (P<0.05).

**Table 3. Branched chain amino acids ratio to lysine in different raw materials.**

Cereals	Amino acids in CP (%)				
	Ile	Lys	Leu	Val	Ile:Leu:Val:Lys
Barley	3.44	3.66	6.79	4.83	94:185:132:100
Corn	3.38	3.10	11.89	4.64	109:383:149:100
Wheat	3.35	2.86	6.61	4.17	117:231:146:100
Rapeseed meal	3.90	5.13	6.82	5.12	76:133:99:100
Sunflower meal	4.01	3.48	6.22	5.03	115:179:144:100
Soybean meal (48%)	4.55	6.04	7.59	4.73	75:126:78:100
Blood plasma	3.36	8.75	9.56	4.73	38:109:54:100

## Low crude protein

There are four major reasons for reducing the CP in an intensive farming system:

- Sparing expensive resources like protein sources.
- Reducing feed cost without losing the animal performance.
- Compliance with the nitrogen excretion regulation.
- Improving the long-term sustainability of animal production.

Until now, commercial nutritionists were unable to go below a certain level of the CP because of the lack of Ile as a supplementary source. Nowadays, Ile can be very well incorporated in formulating low CP diets without compromising Ile requirements under different raw materials regimes.

Recently, Van Harn et al., (2019) tested the lowering of CP in grower (CP:20.8-17.8%) and finisher (CP:19.8-16.8%) diets in broilers. The diet was based on wheat, soybean meal (SBM), rapeseed meal (RSM) and corn, which was fortified with the crystalline amino acids to meet their requirements in low CP treatments (Table 4).

Continuous reduction in CP from 19.1 to 16.8% while keeping the nutrient contents of the diet constant did not impact the performance of the animal (Table 5). However, there was a significant improvement in FCR in the diet with 17.6% CP as compared to the control diet (CP:19.1%).

*Continued on page 28*

Items	Dietary composition					Nutrient composition			
	Control	CP-1%	CP-2%	CP-3%		Control	CP-1%	CP-2%	CP-3%
Wheat	39.64	30	30	30.29	AME Kcal/kg	3025	3025	3025	3025
Corn	20.56	33.02	36.49	40	CP% (Analysed)	191	18.4	17.6	16.8
SBM (48%)	24.99	22.55	18.41	14.62	Lys	9.9	9.9	9.9	9.9
RSM	4.28	4.4	5	5	M+C	7.5	7.5	7.5	7.5
MCP	0.15	0.19	0.22	0.26	Thr	6.4	6.4	6.4	6.4
Lys	0.16	0.24	0.36	0.47	Trp	2.17	2.17	2.17	2.17
Met	0.21	0.24	0.27	0.31	Ile	7	7	7	7
Thr	0.05	0.08	0.14	0.19	Val	7.9	7.9	7.9	7.9
Val	0.02	0.07	0.14	0.2	Arg	11.3	11.3	11.3	11.3
Arg		0.08	0.19	0.3	Gly+Ser	2.8	2.8	2.8	2.8
Ile		0.04	0.11	0.18		Ideal amino acid of BCAA			
Gly		0.09	0.04	0.06	Val:Lys	0.80	0.80	0.80	0.80
Trp		0.02	0.04	0.06	Ile:Lys	0.71	0.71	0.71	0.71

**Table 4. Composition of the experimental diets.**

Continued from page 27

### Conclusion

L-isoleucine is an essential amino acid in broilers. The ideal dlle:dLys in broilers is 0.67, 0.68 and 0.69 for starter, grower and finisher periods, respectively. These ratios are a little higher than what is currently being achieved in industrial formulations.

However, a constant dlle:dLys ratio of 0.69:1.00 is suggested in all age periods in order to compensate batch to batch variations in protein sources under commercial conditions.

In combination with other amino acids, Ile supports the formulation of economical feed when incorporated in a low CP diet without losing animal performance. ■

**Table 5. Performance parameters 0-35 days of age in broilers.**

Crude protein % (calculated)	0-35 days				P value
	19.1 (19.8)	18.4 (18.8)	17.6 (17.8)	16.8 (16.8)	
Live weight (g)	2416	2431	2447	2448	0.595
BWG (g/d)	68.0	68.4	68.8	68.9	0.595
FCR (g/g)	1.549 <sup>a</sup>	1.542 <sup>ab</sup>	1.505 <sup>c</sup>	1.510 <sup>bc</sup>	0.002
Feed intake (g/d)	105.2	105.4	103.6	103.9	0.147